


SURVEY HIGHLIGHTS ON THE VULNERABILITY OF ENGLISH-SPEAKING FATHERS




High Psychological Distress

English-speaking Respondents




French-speaking Counterparts




Suicidal Ideation

English-speaking Respondents




French-speaking Counterparts




Have seriously considered suicide in the last 12 months

Adjusting to Fatherhood




Found it **difficult (very or somewhat)** to adjust to fatherhood



Psychosocial Consultation


English-speaking Respondents



Consulted a psychosocial resource or practitioner in the last year

Violence in the Family Environment

English-speaking respondents who have experienced **at least one form of violence** in their family environment **often/ occasionally/ rarely**




Form of Violence	English-speaking Respondents	French-speaking Counterparts
Psychological abuse	42%	34%
Severe physical violence	31%	16%
Sexual abuse	16%	6%


Medical Consultation

English-speaking Respondents

Have consulted a doctor or other health care professional in the 12 months prior to the survey



French-speaking Counterparts



Secrétariat aux relations avec les Québécois d'expression anglaise



These results are part of a [larger study](#) realized by the Regroupement pour la valorisation de la paternité in collaboration with SOM, survey done with the financial contribution of the Secrétariat aux relations avec les Québécois d'expression anglaise.

Source: SOM, 2022. Survey on Vulnerability in the Context of Fatherhood: Report.